

Restraints in Phenix

Generating & modifying for all scenarios

Nigel W. Moriarty

User's Meeting
University of Kansas, November 2023

What are restraints?

- Restraints are harmonic functions that provide (mostly) chemical information via residuals & gradients
- Needed because experimental data is rarely sufficient for structure determination
- Weighted with the experimental information to find the optimal result


Restraints in Action

- Libraries
 - Monomer Library
 - GeoStd
- Algorithms
 - Polymer
 - Links

GeoStd

- All standard amino acids
- Current list of non-standard amino acids
- All standard RNA/DNA
- Current list of non-standard RNA/DNA
- Others

What you will see



Or

Sorry: Fatal problems interpreting model file:

Number of atoms with unknown nonbonded energy type symbols: 21

Please edit the model file to resolve the problems and/or supply a CIF file with matching restraint definitions, along with `apply_cif_modification` and `apply_cif_link` parameter definitions if necessary.

CIF

- Crystallographic Information File
- mmCIF – macro-molecular CIF
- Used for
 - Model
 - Data
 - Maps
 - Ligands
 - Information
 - Restraints

Confusion

- All depositions of X-ray model use mmCIF from 1 July 2019
- “I need a CIF file.”
 - But what do you really need?


Restraints?

- Provide a reasonable geometry during refinement particularly at low resolution
 - Bonds, angles, dihedrals, chirals, planes, ...
- Must be weighted against the experimental information

Overview

- eLBOW - electronic Ligand Builder & Optimisation Workbench
- ReadySet! - One-stop preparation for your refinement needs
- REEL - Restraints Editor Essentially Ligands


Ligands in crystallography



eLBOW goals

- Fast, simple and flexible procedure to include ligands
- Reduce the tedium of building 3D ligand models
- Automate generation of restraints for ligands
- Comparison of ligand structures

Chemical input




Chemical restraints (CIF)
Cartesian coordinates (PDB)

Reflection data
Protein information

refinement

N. W. Moriarty, R. W. Grosse-Kunstleve, P. D. Adams, (2009). Acta Cryst. D 65, 1074-1080.

Topology



Optimisation

Topology information - Atoms, bonds, angles, ...



Construct Z-Matrix



Simple force field geometry optimisation



Add hydrogens



AM1 geometry optimisation




Output geometry (PDB) and restraints (CIF)

Getting ready to refine

- Many details needed to prepare for structure refinement

Chemical input



Chemical restraints (CIF)
Cartesian coordinates (PDB)

Protein Information

ReadySet!

Experimental data


refinement

ReadySet!

- Add hydrogens
 - Default: adds hydrogens to protein, ligands
 - Protein - Reduce
 - Ligands - eLBOW
 - Add hydrogens to water
 - Add deuteriums instead of hydrogens
 - Add hydrogen & deuteriums appropriately
- Generate restraints



ReadySet!



ReadySet!

- Restraints CIF filename
- Restraints CIF directory
- LINKS to “edits”
- --dry-run to show ligand process pathway
- Metal coordination


Ligand Fitting




Approach

- Where is the ligand?
 - Choose the largest region of contiguous density
- What are rotatable bonds?
 - Analyze ligand for allowed rotations
- What is the orientation of the ligand?
 - Fit core of ligand
- What is the conformation of the ligand?
 - Trace the ligand out from the core

Automated Ligand Fitting



The best fit is measured using the correlation of the model to the electron density



... and extend the model into the density




FFT methods are used to rapidly fit fragments to density

The extension uses rotation around torsion angles to find the best fit

Choose largest rigid fragments of ligand



... and identify their best locations

Take each best location...





Fitting Over a Range of Resolutions

FAD (0.95 Å, 1NIP)



8-(2,5-DIMETHOXY-BENZYL)-2-FLUORO-9-PENT-9H-PURIN-6-YLAMINE (2.2 Å 1UYI)

ATP (3 Å, 1NBM)



(1-(4-IODOBENZOYL)-5-METHOXY-2-METHYL-INDOLE-3-ACETIC ACID (4.5 Å, 1PGF)

Results

- Based on 6209 PDB entries with 9319 ligands

	Ligands in PDB fitting Fo-Fc map with CC ≥ 0.75	All Ligands
Number of ligands	6590	9319
Mean CC of ligand in PDB	0.85	0.78
Mean CC of fitted ligand (unrefined)	0.76	0.72
Rmsd ≤ 1.0 Å	2715 (41%)	3004 (32%)
Rmsd ≤ 2.0 Å	4666 (71%)	5421 (58%)
Rmsd > 10 Å	310 (5%)	1417 (15%)

Restraints in phenix.(real_space_)refine

- LINK records have no impact
- Automatically accesses the “standard” residues restraints
- Automatically links the “standard” residues
- Parameter “link_all=True” links
 - Covalent ligands
 - Carbohydrates
 - Metal ions

phenix.(real_space_)refine (continued)

- RNA/DNA restraints
 - Base pair hydrogen bonding
 - Base pair planarity
 - Base stacking (parallelity)
- Secondary Structure restraints
- NCS restraints
- Custom bonds & angles using edits
- Restraints are written to .geo file including non bonded interactions


QM Restraints

- Generates restraints of ligands using Quantum Mechanics *in situ*
- There are two ways of using QMR
 - In *phenix.refine*
 - In a standalone program *mmtbx.quantum_interface*
- Python2 needs \$PHENIX_MOPAC
- For ORCA, set \$PHENIX_ORCA

QMR

In situ restraints generation

- Carve out the ligand environment
- Minimise the ligand geometry *in situ*
- Transfer geometry values to restraints (and write to disk)
- Refinement with modified restraints



QI

- `phenix.fetch_pdb 4W53 --mtz`
- `phenix.ready_set 4W53.pdb`
- `mmtbx.quantum_interface 4W53.updated.pdb
format=qi write_qmr_phil=True`

The result is a PHIL scope for MBN QMR restraints:

- `4W53.updated_A_200_MBN.phil`

QMR phil

```
qi.qm_restraints {  
  selection = "chain A and resid 200 and rename MBN"  
  run_in_macro_cycles = *first_only first_and_last all last_only test  
  buffer = 3.5
```

```
  calculate = *in_situ_opt starting_energy final_energy starting_strain final_strain starting_bound  
  final_bound starting_higher_single_point final_higher_single_point
```

```
  write_files = *restraints pdb_core pdb_buffer pdb_final_core *pdb_final_buffer
```

```
package {  
  program = *mopac test  
  charge = Auto  
  multiplicity = Auto  
  method = Auto  
  basis_set = Auto  
  solvent_model = None  
  nproc = 1  
  read_output_to_skip_opt_if_available = True  
  ignore_input_differences = False  
  view_output = None}}
```

Run

- mmtbx.quantum_interface 4W53.updated.pdb 4W53.updated_A_200_MBN.phil
run_qmr=True

QM energies


```
"chain A and resid 200 and resname MBN"
```

```
Macro cycle 1
```

```
strain                2.073 kcal/mol (atoms 15)
```

```
opt                   -405946.999 kcal/mol (atoms 246)
```

In situ



Summary

- eLBOW & ReadySet! perform better when provided with better input. (GIGO)
- Need to know something about the ligand
 - Hierarchy of input file value
- Check your .geo file for confirmation of restraints